

Form PTO-1390  
(REV 10-2000)

U S DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

1581.0800000

U.S. APPLICATION NO. (IF KNOWN, SEE 37 C.F.R. § 1.5)

N/A 09/831050

INTERNATIONAL APPLICATION NO

PCT/GB99/03699

INTERNATIONAL FILING DATE

5 NOVEMBER 1999

PRIORITY DATE CLAIMED

5 NOVEMBER 1998

TITLE OF INVENTION

DELIVERY OF SUPEROXIDE DISMUTASE TO NEURONAL CELLS

APPLICANT(S) FOR DO/EO/US

SHONE *et al.*

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)).
4. ☒ The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ has been communicated by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ have been communicated by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 372(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 C.F.R. 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.  
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Authorization To Treat A Reply As Incorporating An Extension Of Time Under 37 C.F.R. § 1.136(a)(3) (*in duplicate*);  
Copy of the International Preliminary Examination Report with Annexes;  
Copy of the International Search Report;  
Application Data Sheet; and  
Two (2) return postcards.

U.S. APPLICATION NO. (if known, see 37 CFR 1.50) N/A <b>097831050</b>		INTERNATIONAL APPLICATION NO PCT/GB99/03699		ATTORNEY'S DOCKET NUMBER 1581.0800000	
--	--	--	--	--	--

17. <input checked="" type="checkbox"/> The following fees are submitted:	CALCULATIONS      PTO USE ONLY
<b>Basic National Fee (37 CFR 1.492(a)(1)-(5)):</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... <b>\$1000.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... <b>\$860.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... <b>\$710.00</b>  International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... <b>\$690.00</b>  International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4) ..... <b>\$ 100.00</b>	                      <b>\$860.00</b>
<b>ENTER APPROPRIATE BASIC FEE AMOUNT</b> =	<b>\$ 860.00</b>
Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).	<b>\$ 130.00</b>

Claims	Number Filed	Number Extra	Rate			
Total Claims	19 - 20 =	0	X \$18.00	\$ 0.00		
Independent Claims	3 - 3 =	0	X \$80.00	\$ 0.00		
Multiple dependent claim(s) (if applicable)			+ \$270.00	\$		
<b>TOTAL OF ABOVE CALCULATIONS</b>				=	\$ 990.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.					\$	
<b>SUBTOTAL</b>				=	\$ 990.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+	\$	
<b>TOTAL NATIONAL FEE</b>				=	\$ 990.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				+	\$	
<b>TOTAL FEES ENCLOSED</b>				=	\$ 990.00	
					Amount to be refunded:	\$
					charged:	\$


a. ☒ A check in the amount of \$990.00 to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_ to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 19-0036. A duplicate copy of this sheet is enclosed.

**NOTE: Where an appropriate time limit Under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO: <b>STERNE, KESSLER, GOLDSTEIN &amp; FOX P.L.L.C.</b> 1100 New York Avenue, NW, Suite 600 Washington, D.C. 20005-3934	<div style="text-align: center;">             SIGNATURE         </div> <div style="text-align: center;"> <u>Robert W. Esmond</u>            NAME         </div> <div style="text-align: center;"> <u>32,893</u>            REGISTRATION NUMBER         </div>
---	---



PTO/PCT Rec'd 20 AUG 2001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Shone *et al.*

Appl. No. 09/831,050 (U.S. National  
Phase of PCT/GB99/03699)

Int'l Filing Date: November 5, 1999

For: **Delivery of Superoxide Dismutase  
to Neuronal Cells**

Confirmation No.:

Art Unit: *To Be Assigned*

Examiner: *To Be Assigned*

Atty. Docket: 1581.0800000/RWE/M-G

**Second Preliminary Amendment**

Commissioner for Patents  
**Box PCT**  
Washington, D.C. 20231

Sir:

Prior to examination of the above-identified patent application, Applicants respectfully request entry of the following amendments. This Amendment is provided in the following format:

- (A) A clean version of each replacement paragraph/section/claim along with clear instructions for entry;
- (B) Starting on a separate page, appropriate remarks and arguments. 37 C.F.R. § 1.111 and MPEP 714; and
- (C) Starting on a separate page, a marked-up version entitled: "Version with markings to show changes made."

It is not believed that extensions of time or fees for net addition of claims are required beyond those that may otherwise be provided for in documents accompanying this paper. However, if additional extensions of time are necessary to prevent abandonment of this

RECEIVED "0507E860"

application, then such extensions of time are hereby petitioned under 37 C.F.R. § 1.136(a), and any fees required therefor (including fees for net addition of claims) are hereby authorized to be charged to our Deposit Account No. 19-0036.

***Amendments***

***In the Specification:***

Please insert the following section at page 1, after the title of the application:

Cross Reference to Related Application

This is the 371 National Stage of International Application No. PCT/GB99/03699, filed November 5, 1999, published in English on May 18, 2000.

***In the Abstract:***

Please insert the Abstract enclosed on a separate sheet after the claims of the above-captioned application.

09/831,050-0001


**Remarks**

The specification has been amended to add a section entitled "Cross Reference to Related Application" at page 1 after the title, and an abstract on a separate sheet has been added. None of these changes is believed to introduce new matter, and their entry is respectfully requested.

Applicants submit that this application is in condition for examination.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Robert W. Esmond  
Attorney for Applicants  
Registration No. 32,893

Date: Aug 20, 2001

1100 New York Avenue, N.W.  
Suite 600  
Washington, D.C. 20005-3934  
(202) 371-2600

***In the Specification:***

A section entitled "Cross Reference to Related Application" was added at page 1 of the application after the title.

***In the Abstract:***

An abstract on a separate sheet has been added.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Shone *et al.*

Appl. No. (U.S. National Phase of  
PCT/GB99/03699)

Filed: (International Filing Date:  
November 5, 1999)

For: **Delivery of Superoxide Dismutase  
to Neuronal Cells**

Confirmation No.:

Art Unit: *To Be Assigned*

Examiner: *To Be Assigned*

Atty. Docket: 1501.0800000/RWE/M-G

**Preliminary Amendment**

Commissioner for Patents  
Washington, D.C. 20231

Sir:

Before examining the above application, please amend the application as follows. This

Amendment is provided in the following format:

(A) A clean version of each replacement paragraph/section/claim along with clear instructions for entry;

(B) Starting on a separate page, appropriate remarks and arguments. 37 C.F.R. § 1.111 and MPEP 714; and

(C) Starting on a separate page, a marked-up version entitled: "Version with markings to show changes made."

It is not believed that extensions of time or fees for net addition of claims are required beyond those that may otherwise be provided for in documents accompanying this paper. However, if additional extensions of time are necessary to prevent abandonment of this application, then such extensions of time are hereby petitioned under 37 C.F.R. § 1.136(a), and

RECEIVED "050103699"

any fees required therefor (including fees for net addition of claims) are hereby authorized to be charged to our Deposit Account No. 19-0036.

***Amendments***

***In the Claims:***

The following amendments are made with respect to the amended claims annexed to the International Preliminary Examination Report.

Please cancel claims 1-24 without prejudice or disclaimer.

Please add the following claims:

--25. (new) A composition for delivery of superoxide dismutase (SOD) to neuronal cells, comprising SOD linked by a cleavable linker to a neuronal cell targeting component, wherein said neuronal cell targeting component comprises a first domain that binds to a neuronal cell and a second domain that translocates the SOD of the composition into the neuronal cell.

26. (new) The composition of claim 25 for delivery of SOD to mitochondria of neuronal cells wherein the SOD comprises a sequence targeting the SOD to mitochondria in the neuronal cell.

27. (new) The composition of claim 26 wherein the SOD is a hybrid of Mn-SOD and a sequence targeting the hybrid to mitochondria.



[illegible][illegible][illegible]

- [illegible]

**Figure 1**

Figure 1 shows the results of the first two steps of the hierarchical regression analysis. The dependent variable is the change in perceived stress from baseline to follow-up. The independent variables are age, gender, education, income, and marital status. The model explains 10% of the variance in the change in perceived stress.

Variable	B	SE	t	p
Age	-0.001	0.001	-0.1	.921
Gender	0.001	0.001	0.1	.921
Education	0.001	0.001	0.1	.921
Income	0.001	0.001	0.1	.921
Marital Status	0.001	0.001	0.1	.921

- [illegible]

[illegible][illegible]

34. (new) A method of delivering SOD to a neuronal cell comprising administering to a mammal an effective amount of the composition of claim 33.

35. (new) The method of claim 34 comprising injecting the composition.

36. (new) A composition for delivery of a therapeutic agent to neuronal cells, comprising the therapeutic agent linked by a cleavable linker to a neuronal cell targeting component, wherein said neuronal cell targeting component comprises a first domain that binds to a neuronal cell and a second domain that translocates the therapeutic agent of the composition into the neuronal cell.

37. (new) A polypeptide comprising a bacterial SOD or derivative thereof and a sequence for targeting the polypeptide to a human mitochondria.

38. (new) The polypeptide of claim 37 wherein the SOD is from *Bacillus*.

39. (new) The polypeptide of claim 37 which is a fusion protein.

40. (new) A nucleotide encoding the polypeptide of claim 37.

41. (new) A cell comprising the nucleotide sequence of claim 40.

42. (new) The composition of claim 25 wherein the cleavable linker is a disulphide bridge between first and second cysteine residues, wherein said first cysteine residue is on the SOD and said second cysteine residue is on the neuronal cell targeting component.

43. (new) The composition of claim 25 wherein the cleavable linker is a site for a protease found in neuronal cells.

TECHNICAL STAFF

**Remarks**

Applicants respectfully request that this Preliminary Amendment be entered by the Examiner. By the foregoing amendment, claims 1-24 are sought to be cancelled without prejudice to or disclaimer of the subject matter therein. New claims 25-43 are sought to be added. These changes are believed to introduce no new matter, and their entry is respectfully requested. Support for the new claims can be found in the specification and original claims.


Upon entry of the foregoing amendment, claims 25-43 are pending in the application, with claims 25, 36 and 37 being the independent claims.

Applicants respectfully request that this Preliminary Amendment under 37 C.F.R. § 1.111 be entered by the Examiner. If the Examiner believes that personal communication will expedite prosecution of this application, the Examiner is invited to telephone the undersigned at the number provided.

Prompt and favorable consideration of this application is respectfully requested.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Robert W. Esmond  
Attorney for Applicants  
Registration No. 32,893

Date: May 4, 2001

1100 New York Avenue, N.W.  
Suite 600  
Washington, D.C. 20005-3934  
(202) 371-2600

**Version with markings to show changes made**

***In the claims***

Please cancel claims 1-24.

Please add claims 25-43.

T00220"030T0300

- 54 -

**CLAIMS**

1. A composition for delivery of superoxide dismutase (SOD) to neuronal cells, comprising:-

5 SOD; linked by a cleavable linker to  
a neuronal cell targeting component, comprising a first domain that binds to a neuronal cell and a second domain that translocates the SOD of the composition into the neuronal cell, wherein, after translocation of the SOD into the cell, the linker is cleaved to release  
10 SOD from the neuronal cell targeting component.

2. A composition according to Claim 1 wherein the cleavable linker is:-

- 15 - a disulphide bridge between cysteine residues, one residue on the SOD and one residue on the neuronal cell targeting component; or  
- a site for a protease found in neuronal cells.

3. A composition according to Claim 1 or 2 for delivery of SOD to mitochondria of neuronal cells wherein the SOD comprises a sequence  
20 targeting the SOD to mitochondria in the neuronal cell.

4. A composition according to Claim 3 wherein the SOD is a hybrid of Mn-SOD and a sequence targeting the hybrid to mitochondria.

25 5. A composition according to Claim 3 or 4 wherein the mitochondria targeting sequence is derived from human Mn-SOD.

30 6. A composition according to any of Claims 1-5 wherein the SOD is bacterial SOD or is a derivative thereof that substantially retains the superoxide dismutase activity of bacterial SOD.

- 55 -

7. A composition according to any of Claims 1 to 6 wherein the first domain is selected from (a) neuronal cell binding domains of clostridial toxins; and (b) fragments, variants and derivatives of the domains in (a) that substantially retain the neuronal cell binding activity of the domains of (a).

8. A composition according to any Claims 1 to 7 wherein the second domain is selected from (a) domains of clostridial neurotoxins that translocate polypeptide sequences into cells, and (b) fragments, variants and derivatives of the domains of (a) that substantially retain the translocating activity of the domains of (a).

9. A composition according to any of Claims 1 to 8 wherein the linker is a disulphide bridge.

10. A pharmaceutical composition for treatment of oxidative damage to neuronal cells comprising a composition according to any of Claims 1 to 9 and a pharmaceutically acceptable carrier.

11. A method of delivering SOD to a neuronal cell comprising administering a composition according to Claim 10.

12. A method according to Claim 11 comprising injecting the composition.

13. A method of making a composition according to any of Claims 1 to 8 comprising chemically linking SOD, a linker and a neuronal cell targeting component.

14. A method of making a composition according to any of Claims 1 to 9 comprising expressing a DNA that codes for a polypeptide having SOD activity, a linker, and a neuronal cell targeting component.

- 56 -

15. A method according to claim 14 wherein the polypeptide further comprises a purification sequence and the method further comprises purifying the polypeptide and then cleaving the polypeptide to remove the purification sequence to leave SOD, the linker and the neuronal cell targeting component.

16. A composition for delivery of a therapeutic agent to neuronal cells, comprising:-

the therapeutic agent; linked by a cleavable linker to a neuronal cell targeting component, comprising a first domain that binds to a neuronal cell and a second domain that translocates the therapeutic agent of the composition into the neuronal cell wherein, after translocation of the SOD into the cell, the linker is cleaved to release SOD from the neuronal cell targeting component.

17. A composition for delivery of a therapeutic agent to neuronal cells according to Claim 16, wherein the cleavable linker is either a disulphide bridge or a site for a protease found in neuronal cells.

18. A polypeptide comprising a bacterial SOD or derivative thereof that substantially retains the superoxide dismutase activity of bacterial SOD and a sequence for targeting the polypeptide to a human mitochondria.

19. A polypeptide according to Claim 18 wherein the SOD is from *Bacillus*.

20. A polypeptide according to Claim 18 or 19 which is a fusion protein.

21. A nucleotide encoding the polypeptide of any of Claims 18-20.

22. A vector comprising the nucleotide of Claim 21.



- 57 -

23. A method of making a polypeptide according to any of Claims 18-20 comprising expressing the nucleotide sequence of Claim 21.

24. A cell comprising the nucleotide sequence of Claim 21 or the vector of Claim 22.

5

4/PK TS

- 1 -

DELIVERY OF SUPEROXIDE DISMUTASE TO NEURONAL CELLS

5 The present invention relates to compositions and methods for delivery of superoxide dismutase (SOD) to neuronal cells, and in particular for delivery of SOD to mitochondria in those cells. The present invention also provides methods of making the constructs.

10 A number of nervous system disorders would benefit considerably from rapid intervention with several therapeutic agents. Examples of such disorders are global, focal or spinal cord ischaemia induced by stroke or injury. Neurones injured by trauma or ischaemia produce elevated levels of transmitter substances which result in high levels of reactive oxygen species. These reactive oxygen species, in high concentrations are toxic to both the neurones and the surrounding cells which potentiates and amplifies the damage process. Rapid therapeutic intervention with agents to reduce oxidative stress in cases of neuronal injury caused by stroke or trauma could therefore significantly limit this secondary damage process.

15 One such potential therapeutic agent is superoxide dismutase which neutralises the harmful effects of the superoxide radicals by converting them to hydrogen peroxide and oxygen.

20

25 Francis and co-workers demonstrated that post-ischaemic infusion of Copper/Zinc superoxide dismutase (Cu/Zn-SOD) reduces cerebral infarction following ischaemia/reperfusion in rats (Experimental Neurology (1997) 146, 435-443) through the reduction of damaging free-radical oxygen. Lim et al. have shown that administration of Cu/Zn-SOD attenuates the level of reperfusion injury following spinal cord ischaemia in dogs (Ann. Thorac. Surg. (1986) 42, 282-286). Cuevas et al. have similarly demonstrated protective effects of SOD, both on neurological recovery and spinal infarction, in ischaemic reperfusion injury of the rabbit spinal cord (Acta Anat. (1990) 137, 303-310. A major problem in the use of such therapies is the maintenance of useful concentration of the active agent at the site

30

- 2 -

of trauma. Enzymes such as Cu/Zn-SOD are rapidly cleared from the systemic circulation; in the case of the latter enzyme the  $t_{1/2}$  in rat is 4-8 minutes. A number of strategies have been employed to overcome these difficulties. Matsumiya et al. (Stroke (1991) 22, 1193-1200) conjugated Cu/Zn-SOD to polyethylene glycol to increase its half-life in the blood. Francis and co-workers (Experimental Neurology (1997) 146, 435-443) describe the neuronal delivery of Cu/Zn-SOD by fusion of the enzyme to the binding domain of tetanus toxin. None of these strategies, however, are particularly efficient at delivering the enzyme to the intracellular neuronal compartments where the enzyme can be effective.

In most eukaryotic species, two intracellular forms of superoxide dismutase exist: the Cu/Zn-SOD which is located within the cytoplasmic and nuclear compartments and manganese superoxide dismutase (Mn-SOD) which is located within the mitochondrial matrix. Human Mn-SOD is a tetrameric enzyme and is larger than the dimeric Cu/Zn-SOD. Several studies have shown that decreased Mn-SOD may be associated with one or more chronic diseases such as ovarian cancer (Nishida *et al.* (1995) Oncology Reports, 2, 643-646) and diabetes (L'Abbe *et al.*, (1994) Proc Soc Exp Biol Med, 207, 206-274). In addition, mice in which the Mn-SOD gene has been knocked out exhibit several novel pathogenic phenotypes including severe anaemia, degeneration of neurones in the basal ganglia and brainstem, and progressive motor disturbances characterised by weakness and rapid fatigue (Lebovitz *et al.*, (1996) Proc Natl Acad Sci USA, 93, 9782-9787). In addition these mice showed extensive damage to the neuronal mitochondria. Overexpression of Mn-SOD in cell lines and transgenic mice showed that damage and apoptosis of neurones under oxidative stress was markedly reduced (Keller *et al.*, (1998) Journal of Neuroscience, 18, 687-697). Mitochondrial damage was also reduced. These data showed that superoxide accumulation and subsequent mitochondrial damage play key roles in neuronal death induced by trauma both *in vitro* and *in vivo*. Delivery of agents which reduce the level of

- 3 -

oxidative stress to neuronal cells may therefore reduce neuronal cell death and afford considerable therapeutic benefits.

5 Mn-SODs of bacterial origin such as that from *Bacillus stearothermophilus* or *B. caldotenax* consist of two subunits and are smaller than the human isoform which is a tetramer. *B. stearothermophilus* and *B. caldotenax* Mn-SOD also have considerably lower immunogenicity than the human isoform which is an advantage for continued therapeutic use. As enzymes for  
10 therapeutic applications, however, they suffer from similar drawbacks to other SODs in that very little of the administered enzyme is retained within the tissues where it would be therapeutically beneficial.

15 The botulinum neurotoxins are a family of seven structurally similar, yet antigenically different, protein toxins whose primary site of action is the neuromuscular junction where they block the release of the transmitter acetylcholine. The action of these toxins on the peripheral nervous system of man and animals results in the syndrome botulism, which is characterised by widespread flaccid muscular paralysis (Shone (1986) in 'Natural Toxicants in Foods', Editor D. Watson, Ellis Harwood, UK). Each  
20 of the botulinum neurotoxins consist of two disulphide-linked subunits; a 100 kDa heavy subunit which plays a role in the initial binding and internalisation of the neurotoxin into the nerve ending (Dolly et. al. (1984) Nature, 307, 457-460) and a 50 kDa light subunit which acts intracellularly to block the exocytosis process (McInnes and Dolly (1990) Febs Lett., 261, 323-326; de Paiva and Dolly (1990) Febs Lett., 277, 171-174). Thus it is  
25 the heavy chains of the botulinum neurotoxins that impart their remarkable neuronal specificity.

30 Tetanus toxin is structurally very similar to botulinum neurotoxins but its primary site of action is the central nervous system where it blocks the release of inhibitory neurotransmitters from central synapses (Renshaw cells). As described for the botulinum toxins above, it is domains within

- 4 -

the heavy chain of tetanus toxin that bind to receptors on neuronal cells.

5 The binding and internalisation (translocation) functions of the clostridial neurotoxin (tetanus and botulinum) heavy chains can be assigned to at least two domains within their structures. The initial binding step is energy-independent and appears to be mediated by one or more domains within the H<sub>C</sub> fragment of the neurotoxin (C-terminal fragment of approximately 50kDa) (Shone *et al.* (1985), Eur. J. Biochem., 151, 75-82) while the translocation step is energy-dependent and appears to be mediated by one or more domains within the H<sub>N</sub> fragment of the neurotoxin (N-terminal fragment of approximately 50kDa).

10 Isolated heavy chains are non-toxic compared to the native neurotoxins and yet retain the high affinity binding for neuronal cells. Tetanus and the botulinum neurotoxins from most of the seven serotypes, together with their derived heavy chains, have been shown to bind a wide variety of neuronal cell types with high affinities in the nM range (e.g botulinum type B neurotoxin; Evans *et al.* (1986) Eur. J. Biochem. 154, 409-416).

20 Another key characteristic of the binding of these neurotoxins is that tetanus, botulinum A, B, C<sub>1</sub>, D, E and F neurotoxins all appear to recognise distinct receptor populations, and collectively the clostridial neurotoxin heavy chains provide high affinity binding ligands that recognise a whole family of receptors that are specific to neuronal cells.

25 However, whilst it is known to provide a fusion of a SOD with a neurotoxin heavy chain, this fusion has been found to be ineffective for delivery of SOD to neuronal cells and inactive in *in vitro* assays for potential therapeutic activity.

30 It is an object of the invention to provide compositions and methods for delivery of SOD to neuronal cells. A further object is to provide

- 5 -

compositions and methods for treatment or at least amelioration of conditions such as ischemic stroke.

Accordingly, the present invention provides a construct for delivering superoxide dismutase (SOD) to neuronal cells. In one aspect of the invention the construct consists of a SOD which has been combined with various functional protein domains to effect efficient targeting to the mitochondria within neuronal cells. The construct of specific embodiments of the invention, described in further detail below, contains the following elements:-

- a SOD which contains a leader sequence for targeting SOD to the mitochondria;
- a dimeric SOD which has low immunogenicity and high stability;
- a SOD which is linked to a domain that effects translocation across lipid membranes;
- a linkage between the SOD and the translocation domain that is cleaved within the neuronal cytosol; and
- a domain which selectively targets the construct to neuronal cells.

A first aspect of the invention thus provides a composition for delivery of superoxide dismutase (SOD) to neuronal cells, comprising:-

SOD; linked by a cleavable linker to  
a neuronal cell targeting component, comprising a first domain that binds to a neuronal cell and a second domain that translocates the SOD of the composition into the neuronal cell.

The linker is cleavable and thus, in use, after translocation of the SOD into the cell, the linker is cleaved to release SOD from the neuronal cell targeting domain. A suitable linker is a disulphide bridge between cysteine residues, one residue on the SOD and one residue on the neuronal cell targeting component, for example on the second domain. Another example of a linker is a site for a protease found in neuronal cells. In this way, the

- 6 -

linker is such that it is cleaved within the cell, separating the SOD from the other translocated portions of the composition.

By SOD is meant a sequence that has superoxide dismutase activity, and may also comprise a further sequence or sequences conferring additional properties on that portion of the constructs. For example, the SOD optionally also includes a sequence targeting the SOD to mitochondria in a neuronal cell.

The SOD may be a hybrid of Mn-SOD and a sequence targeting the hybrid to mitochondria. The SOD may be of bacterial or human origin, or a derivative thereof, and may be comprised of sequences from more than one origin, provided that it has superoxide dismutase activity.

The first domain may suitably be selected from (a) neuronal cell binding domains of clostridial toxins; and (b) fragments, variants and derivatives of the domains in (a) that substantially retain the neuronal cell binding activity of the domains of (a). The second domain is suitably selected from (a) domains of clostridial neurotoxins that translocate polypeptide sequences into cells, and (b) fragments, variants and derivatives of the domains of (a) that substantially retain the translocating activity of the domains of (a).

In an embodiment of the invention a construct comprises SOD linked by a disulphide bridge to a neuronal cell targetting component comprising a first domain that binds to a neuronal cell and a second domain that translocates the SOD into the neuronal cell. This construct is made recombinantly as a single polypeptide having a cysteine residue on the SOD which forms a disulphide bridge with a cysteine residue on the second domain. The SOD is covalently linked, initially, to the second domain. Following expression of this single polypeptide SOD is cleaved from the second domain leaving the SOD linked only by the disulphide bridge to the rest of the construct.

- 7 -

5 A second aspect of the invention provides a pharmaceutical composition for treatment of oxidative damage to neuronal cells comprising a composition according to the invention with a pharmaceutically acceptable carrier. This composition may be used to deliver SOD to a neuronal cell, for example by administration of the composition by injection.

10 A third aspect of the invention provides a method of preparing a composition according to the invention, comprising chemically linking SOD, linker and neuronal cell targeting components. The SOD preferably is free of cysteine residues and the method preferably comprises treating the SOD with a cross-linker which will form a disulphide bridge with a cysteine residue on the neuronal cell targeting domain.

15 In a further embodiment of the third aspect of the invention, there is provided a method of making a composition according to the invention comprising expressing a DNA that codes for a polypeptide having SOD, a linker, a neuronal cell targeting component. The polypeptide may further comprise a purification sequence and the method may further comprise purifying the polypeptide using this sequence and then cleaving the  
20 polypeptide to remove the purification sequence to leave SOD, the linker and the neuronal cell targeting component.

25 By virtue of the combination of properties defined above, constructs of the invention are surprisingly efficient at transporting SOD to the mitochondria within neuronal cells. The ability of the superoxide dismutase to be translocated into the cytosol by virtue of the 'translocation domain' within the construct and the cleavage of the enzyme from the latter domain within the cell is key to this targeting efficiency. As such the construct of the invention has considerable therapeutic value in treating neuronal diseases  
30 which results from oxidative stress and has several advantages over previously described SOD formulations. Mitochondria within cells containing high levels of superoxide radicals are particularly sensitive to



- 8 -

damage and the ability of the construct of the invention to target the SOD to these organelles offers considerable advantage in that the enzyme can act to remove the superoxide radicals where it is most required.

5 The construct of the invention may be used clinically in a variety of neuronal diseases which are caused or augmented by oxidative stress. Such conditions include ischaemic stroke, Parkinson's disease, Huntington's disease and motor neurone diseases. In the case of ischaemia/reperfusion injury caused by stroke or trauma, delivery of the construct of the invention to neurones of the hippocampus may afford considerable therapeutic benefits by reducing neuronal damage and death. Other neuronal diseases where the underlying cause is oxidative stress would also benefit from the therapeutic effects of the construct of the invention.

10  
15 In a preferred aspect of the invention, the SOD is a dimeric, manganese superoxide dismutase (Mn-SOD) which is of bacterial origin and has low immunogenicity and high stability.

20 The use of a bacterial Mn-SOD in constructs has a number of advantages compared to the use of the human Mn-SOD isoform:-

25 - the low immunogenicity of the bacterial Mn-SOD is advantageous where repeated administration of the construct is required, in which cases the induction of adverse host immune responses is reduced; and

30 - the smaller size of constructs based on the dimeric bacterial Mn-SOD compared to human Mn-SOD (which is a tetramer) both reduces the likelihood of adverse immune responses and increases the rate of diffusion of the construct to its target tissue.

In exercise of an example of the invention, a bacterial Mn-SOD of low

- 9 -

immunogenicity is derived from either *B. stearothermophilus* (sequence as reported by Brock and Walker (1980) Biochemistry, 19, 2873-2882) or *B. caldotenax* (gene and amino acid sequence as defined by Chambers *et al.*, (1992) FEMS Microbiology Letters, 91, 277-284) to which a mitochondrial leader sequence has been fused to the N-terminus of the protein by recombinant technology. This Mn-SOD-leader hybrid is linked by a disulphide bridge to a translocation domain derived from a bacterial protein toxin, such as botulinum neurotoxin. The translocation domain, in turn is fused to a receptor binding domain derived from a clostridial neurotoxin (botulinum or tetanus). The construct is produced initially as a single polypeptide by recombinant technology and subsequently converted to the construct of the invention by selective cleavage with a proteolytic enzyme. To produce the construct of the invention, a loop motif containing a unique protease site (e.g amino acid sequences specifically cleaved by proteases such as factor Xa, enterokinase, thrombin) and a cysteine residue is introduced between the C-terminus of the Mn-SOD and the N-terminus of the translocation domain such that a disulphide bridge is formed between the Mn-SOD and the translocation domain. Subsequent cleavage of the protease site generates the active construct. The final construct, when analyzed by sodium dodecyl sulphate polyacrylamide gel electrophoresis in the presence of a reducing agent (e.g. dithiothreitol), dissociates into two major bands, one corresponding to the superoxide dismutase enzyme and a second corresponding to a polypeptide which contains the neuronal binding and translocation domains. In the absence of a reducing agent this dissociation does not occur and the complex is observed as a single major band on the gels.

Modification of Mn-SOD from *B. stearothermophilus* by addition of a mitochondrial targeting sequence offers several advantages over the use of human Mn-SOD which contains its own mitochondrial leader sequence. Firstly, the *B. stearothermophilus* Mn-SOD has a high thermal stability and low immunogenicity which allows administration of several doses of the

- 10 -

enzyme without provoking an immune response from the host that would reduces its efficacy. Secondly, the *B. stearotherophilus* Mn-SOD is a small dimeric enzyme unlike the human Mn-SOD which is a tetramer. Recombinant constructs containing the latter enzyme would therefore have to be considerably larger and more complex in their structure.

In an embodiment of the invention, a DNA encoding a construct of the invention is made up by fusion of following DNA fragments commencing at the 5' end of the gene:-

an oligonucleotide encoding a modified human mitochondrial leader sequence (amino acid sequence: MLSRAVCGTSRQLAPALGYLGSRQ (SEQ ID NO:10) or MLSRAVSGTSRQLAPALGYLGSRQ (SEQ ID NO:11);

an oligonucleotide encoding Mn-SOD from *B. stearotherophilus* (coding for the amino acid sequence as defined in Brock and Walker (1980) Biochemistry, 19, 2873-2882);

an oligonucleotide encoding a linker peptide which contains the thrombin protease cleavage site and a cysteine residue for disulphide bridge formation (peptide sequence: CGLVPAGSGP);

an oligonucleotide encoding a translocation domain derived from a botulinum neurotoxin (e.g. a DNA fragment coding for amino acid residues 449-871 of botulinum type A neurotoxin, or a DNA fragment coding for amino acid residues 441-858 of botulinum type B neurotoxin, or a DNA fragment coding for amino acid residues 440-864 of botulinum type F neurotoxin); and

an oligonucleotide encoding the receptor binding domain of a botulinum neurotoxin or tetanus neurotoxin (e.g. a DNA fragment

- 11 -

coding for amino acid residues 872-1296 of botulinum type A neurotoxin, or a DNA fragment coding for amino acid residues 859-1291 of botulinum type B neurotoxin, or a DNA fragment coding for amino acid residues 865-1278 of botulinum type F neurotoxin, or a DNA fragment coding for amino acid residues 880-1315 of tetanus neurotoxin).

The above DNA fragments may be obtained and constructed by standard recombinant DNA methods. Expression and purification of the assembled construct may be obtained with a variety of suitable expression hosts, e.g. *Escherichia coli*, *Bacillus subtilis*.

The translocation domain and neuronal binding domain of the construct may also be derived from combination of different clostridial neurotoxins. For example, the construct of the invention may contain a translocation domain derived from botulinum type F neurotoxin and a binding domain derived from botulinum type A neurotoxin.

A construct of the invention may be produced using protein chemistry techniques. Mn-SOD derived from *B. stearothermophilus* to which a mitochondrial leader sequence has been fused to the N-terminus of the protein by recombinant technology is modified with a heterobifunctional cross-linking reagent such as N-succinimidyl 3-[2-pyridylidithio] propionate (SPDP). The chemically modified enzyme is then combined to a cell targetting domain which contains the binding and translocation functional domains. The latter may be produced by recombinant technology or purified from the neurotoxins of *Clostridium botulinum* or *Clostridium tetani* by established methods. Chemical coupling of the SPDP-treated Mn-SOD may be accomplished using a free cysteine residue on the polypeptide containing the binding and translocation domains to give a construct of the invention.

- 12 -

Constructs of the invention may be introduced into either neuronal or non-neuronal tissue using methods known in the art. By subsequent specific binding to neuronal cell tissue, the targeted construct will exert its therapeutic effects. Alternatively, the construct may be injected near a site requiring therapeutic intervention, e.g. intrathecal or intracranial injection close to a site of trauma or disease.

The construct of the invention may also be administered with other agents which enhance its delivery to its target tissue. An example of such an agent is one which assists the passage of the construct of the invention through the blood-brain barrier to the central nervous system. The construct of the invention may also be administered in formulations with other therapeutic agents or drugs.

The dosage required for the construct of the invention will depend upon the application and could vary between 1µg/kg to 100mg/kg of body weight. The construct of the invention may be produced as a suspension, emulsion, solution or as a freeze dried powder depending on the application and properties of the release vehicle and its therapeutic contents. The construct of the invention may be resuspended or diluted in a variety of pharmaceutically acceptable liquids depending on the application.

"Clostridial neurotoxin" means a neurotoxin corresponding to tetanus neurotoxin or one of the seven botulinum neurotoxin serotypes (type A, B, C<sub>1</sub>, D, E, F or G).

"Bind" in relation to the clostridial binding fragments, means the interaction between the clostridial fragment and one or more cell surface receptors or markers which results in localisation of the binding fragment or construct in the vicinity of the cell.

"Binding domain" of botulinum or tetanus neurotoxins means a domain of

the toxin which retains the property of being able to bind the receptors on neuronal cells in a similar manner to the intact neurotoxin and encompasses native domains and fragments, variants and derivatives that retain this binding function. This property of the binding domain can be assessed in competitive binding assays. In such assays, radiolabelled neurotoxin (e.g. botulinum type A neurotoxin) is contacted with neuronal cells in the presence of various concentrations of non-radiolabelled fragment representing the 'binding domain' of the neurotoxin. The ligand mixture is incubated with the cells, at low temperature (0-3°C) to prevent ligand internalisation, during which competition between the radiolabelled neurotoxin and non-labelled 'binding domain' fragment may occur. In such assays when the unlabelled ligand used is binding domain of botulinum type A neurotoxin (residues 872-1296), the radiolabelled botulinum type A neurotoxin will be displaced from the neuronal cell receptors as the concentration of its non-labelled 'binding domain' is increased. The competition curve obtained in this case will therefore be representative of the behaviour of a 'binding domain' fragment being able to bind the receptors on neuronal cells in a similar manner to the intact neurotoxin. This property of the binding domain may be used to identify other suitable protein domains which have the desired binding properties. Examples of binding domains derived from clostridial neurotoxins are as follows:-

Botulinum type A neurotoxin	- amino acid residues (872 - 1296)
Botulinum type B neurotoxin	- amino acid residues (859 - 1291)
Botulinum type C neurotoxin	- amino acid residues (867 - 1291)
Botulinum type D neurotoxin	- amino acid residues (863 - 1276)
Botulinum type E neurotoxin	- amino acid residues (846 - 1252)
Botulinum type F neurotoxin	- amino acid residues (865 - 1278)
Botulinum type G neurotoxin	- amino acid residues (864 - 1297)
Tetanus neurotoxin	- amino acid residues (880 - 1315)

"Translocation domain" means a domain or fragment of a protein which effects transport of itself and/or other proteins and substances across a

- 14 -

membrane or lipid bilayer and encompasses native domains and fragments, variants and derivatives that retain this binding function. The latter membrane may be that of an endosome where translocation will occur during the process of receptor-mediated endocytosis. Translocation domains can frequently be identified by the property of being able to form measurable pores in lipid membranes at low pH (Shone *et al.* (1987) Eur J. Biochem. 167, 175-180). The latter property of translocation domains may thus be used to identify other protein domains which could function as the translocation domain within the construct of the invention. Examples of translocation domains derived from bacterial neurotoxins are as follows:-

Botulinum type A neurotoxin	- amino acid residues (449 - 871)
Botulinum type B neurotoxin	- amino acid residues (441 - 858)
Botulinum type C neurotoxin	- amino acid residues (442 - 866)
Botulinum type D neurotoxin	- amino acid residues (446 - 862)
Botulinum type E neurotoxin	- amino acid residues (423 - 845)
Botulinum type F neurotoxin	- amino acid residues (440 - 864)
Botulinum type G neurotoxin	- amino acid residues (442 - 863)
Tetanus neurotoxin	- amino acid residues (458 - 879)

"Translocation" in relation to translocation domain, means the internalisation events which occur after modified clostridial binding fragments bind to the cell surface. These events lead to the transport of substances into the cytosol of neuronal cells.

"Unique protease site" means a protease site incorporated into the construct such that the molecule may be proteolysed at pre-determined sites by a selected protease. The specificity of these proteases is such that cleavage to other parts of the construct does not occur. Examples of unique protease sites are the amino acid sequences cleaved by proteases such as: thrombin, factor Xa, enterokinase.

A fourth aspect of the invention provides a composition for delivery of a

- 15 -

therapeutic agent to neuronal cells, comprising:-

the therapeutic agent; linked by a cleavable linker to  
a neuronal cell targeting component, comprising a first domain that  
binds to a neuronal cell and a second domain that translocates the  
therapeutic agent of the composition into the neuronal cell.

Thus, in use, after translocation of the therapeutic agent into the cell, the  
linker is cleaved to release the therapeutic agent from the neuronal cell  
targeting domain. Other optional and preferred embodiments of the fourth  
aspect of the invention are as for the first-third aspects of the invention.

A fifth aspect of the invention provides a polypeptide comprising a  
bacterial SOD, or derivative thereof, and a sequence for targeting the  
polypeptide to a mitochondria, such as a human mitochondria. The  
polypeptide may be chemically obtained by synthesis of otherwise or may  
be a fusion protein, obtained for example by expression of a nucleotide  
coding for the polypeptide.

The invention hence also provides, in a sixth aspect, a nucleotide encoding  
the polypeptide of the fifth aspect and in a seventh aspect a vector  
comprising the nucleotide of the sixth aspect. Also provided in an eighth  
aspect is a method of making a polypeptide according to the fifth aspect  
comprising expressing the nucleotide sequence of the sixth aspect. In a  
ninth aspect is provided a cell comprising the nucleotide sequence of the  
sixth aspect or the vector of the seventh.

There now follows description of specific embodiments of the invention  
illustrated by drawings in which:-

Fig. 1 shows schematic examples of novel Mn-SODs derived from  
*B. stearothermophilus* and *B. caldotenax*. Two examples of mitochondrial  
leader sequences are shown. In one example, a cysteine residue at position



- 16 -

7 has been mutated to a serine residue. This change enables the production of the construct of the invention without the formation of disulphide bridges in undesirable positions;

5 Figure 2 shows schematic examples of novel Mn-SOD fusion proteins showing the use of peptides and proteins to facilitate purification of the enzyme from the production strain. Various protein and peptide tags (such as histidine-6, S-peptide, maltose-binding protein, calmodulin-binding protein) may be fused to the Mn-SOD to allow rapid purification by affinity chromatography methods. Unique protease sites are incorporated between the purification tag and the Mn-SOD to enable removal of the tag after purification. Protein and peptide tags may be removed by treatment of the fusion protein with the relevant specific protease (e.g. factor Xa, thrombin, enterokinase);

10  
15  
20  
25 Fig. 3 shows a recombinant Mn-SOD construct of the invention. From the N-terminus of the protein, the construct consists of the following components:- (1) a mitochondrial leader (targeting) sequence, (2) a Mn-superoxide dismutase, (3) a loop which contains a unique protease site and which allows disulphide bridge formation, (4) a translocation domain, (5) a neuronal targeting domain. The construct is produced as a single polypeptide; subsequent cleavage with a protease specific for the 'unique protease site' contained within the loop region generates the di-chain construct. Purification tags could added to the constructs as exemplified in Figure 3;

30 Fig. 4 shows the production of a Mn-SOD construct by chemical methods. The method uses a recombinant Mn-SOD, purified as described in Example 1 and coupled to a polypeptide containing the translocation and binding domains as described in Example 4; and

Fig. 5 shows the results of an example to demonstrate the protective

- 17 -

effects of a construct of the invention on NG108 cells subjected to oxidative stress by the addition of 50 $\mu$ M duroquinone for four hours.

The application is also accompanied by a sequence listing in which:-

SEQ ID NO: 1 shows the amino acid sequence of Mn-SOD from *B. caldotenax*;

SEQ ID NO: 2 shows the amino acid sequence of Mn-SOD from *B. stearothermophilus*;

SEQ ID NO: 3 shows the amino acid sequence of a construct of the invention comprising Mn-SOD from *B. stearothermophilus*, a linker that can be cleaved by thrombin, and a heavy chain derived from botulinum neurotoxin serotype A;

SEQ ID NO: 4 shows the amino acid sequence of a construct of the invention comprising Mn-SOD from *B. stearothermophilus*, a linker that can be cleaved by thrombin, and a heavy chain derived from botulinum neurotoxin serotype B;

SEQ ID NO: 5 shows the amino acid sequence of a construct of the invention comprising Mn-SOD from *B. stearothermophilus*, a linker that can be cleaved by thrombin, and a heavy chain derived from botulinum neurotoxin serotype F;

SEQ ID NO: 6 shows the amino acid sequence of a construct of the invention comprising a mitochondrial leader sequence from human Mn-SOD, Mn-SOD from *B. stearothermophilus*, a linker that can be cleaved by thrombin, and a heavy chain derived from botulinum neurotoxin serotype A;

- 18 -

SEQ ID NO: 7 shows the amino acid sequence of a construct of the invention comprising a mitochondrial leader sequence from human Mn-SOD, Mn-SOD from *B. stearothermophilus*, a linker that can be cleaved by thrombin, and a heavy chain derived from botulinum neurotoxin serotype B;

SEQ ID NO: 8 shows the amino acid sequence of a construct of the invention comprising a mitochondrial leader sequence from human Mn-SOD, Mn-SOD from *B. stearothermophilus*, a linker that can be cleaved by thrombin, and a heavy chain derived from botulinum neurotoxin serotype F; and

SEQ ID NO: 9 shows the amino acid sequence for a polypeptide comprising a mitochondrial leader sequence from human Mn-SOD and Mn-SOD from *B. stearothermophilus*;

SEQ ID NO: 10 shows the amino acid sequence of a modified human mitochondrial leader sequence; and

SEQ ID NO: 11 shows an amino acid sequence of a modified human mitochondrial leader sequence.

#### Example 1.

#### **Production and purification of novel *B. stearothermophilus* Mn-SOD containing a mitochondrial leader sequence.**

Standard molecular biology protocols were used for all genetic manipulations (eg. Sambrook *et al.* 1989, Molecular Cloning a Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). A synthetic gene encoding the mitochondrial targeting sequence (amino acids 1-27) of the human Mn-SOD gene was cloned as

- 19 -

an *NdeI* - *Bam* HI fragment into an expression vector so that the transcriptional start corresponds to the ATG codon within the *NdeI* site. The Mn-SOD gene from *B. stearrowthermophilus* or *B. caldotenax* was amplified using PCR to give a *Bgl*III site corresponding to the leucine amino acid at position 5 at the 5' end, and a *Bam*HI site outside the stop codon at the 3' end. This *Bs*III-*Bam*HI fragment was cloned into the expression vector carrying the mitochondrial targeting sequence (digested *Bam*HI) to generate "in-frame" gene fusions. In addition to the wild type mitochondrial targeting sequence, a variant was constructed in which the cysteine at position 7 was changed to serine

The recombinant Mn-SOD expressed in pET28a were produced with amino-terminal histidine (6 His) and T7 peptide tags allowing proteins to be purified by affinity chromatography on either a  $\text{Ni}^{2+}$  charged column or an anti-T7 immunoaffinity column (Smith *et al.* 1988, *Journal of Biological Chemistry*, 263: 7211-7215). Incorporation of a factor Xa protease cleavage between the peptide tag and the expressed Mn-SOD allowed this to be removed after purification. Briefly, cultures of *E.coli* BL21 (DE3) pET28a-Mn-SOD were grown in Terrific broth-kanamycin ( $30 \mu\text{gml}^{-1}$ ) to an  $\text{OD}_{600 \text{ nm}}$  of 2.0, and protein expression was induced by the addition of  $500 \mu\text{M}$  IPTG for approximately 2 h. Cells were lysed by freeze/thaw followed by sonication, lysates cleared by centrifugation and supernatants loaded onto an anion exchange column (MonoQ™ column on a Fast Protein Liquid Chromatography system; Pharmacia Biotech, Uppsala, Sweden). Eluted recombinant Mn-SOD was then desalted and further purified by affinity chromatography on a chelating sepharose column charged with  $\text{Ni}^{2+}$  (Pharmacia Biotech, Uppsala, Sweden). After loading proteins onto the column and subsequent washing, the purified Mn-SOD was eluted with imidazole. All buffers used were as specified by the manufacturer.

A 'maltose binding protein' purification tag was also employed for the purification of some batches of Mn-SOD. The use of this system is

- 20 -

described in detail in New England Biolabs Instruction Manual "Protein Fusion and Purification System" (ver 3.02).

Other tags and protease cleavage site may also be incorporated into the sequence to facilitate purification of Mn-SOD as exemplified in Figure 2.

For purification of a novel *B. stearothermophilus* Mn-SOD which was not conjugated to protein purification tag, the following procedure was used. After harvesting, cells were broken by high pressure homogenisation crude extracts were clarified by centrifugation and batch purified on DE-23 cellulose. The fraction eluted with 0.4M NaCl contained the Mn-SOD. This fraction was then further purified by various chromatographic media using the following sequence:-

DEAE-Sepharose ion exchange chromatography at pH 8.0; elution of the Mn-SOD with a NaCl gradient;

hydroxylapatite chromatography at pH 6.8; elution of Mn-SOD with a phosphate gradient at pH 6.8;

ion exchange chromatography on Q-Sepharose at pH 7.5; elution with a NaCl gradient; and

gel filtration on Sephacryl S-200.

The purified Mn-SOD may be dialysed against Hepes buffer (0.1M, pH7.4) containing 0.15M NaCl and stored at -80°C.

#### Example 2.

Preparation and purification of a recombinant Mn-SOD construct of the invention.

- 21 -

Standard molecular biology protocols were used for all genetic manipulations (*eg.* Sambrook *et al.* 1989, Molecular Cloning a Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). Various DNA fragments of the construct were generated using Recursive PCR reactions (Prodromou & Pearl 1992, *Protein Engineering*, 5: 827-829) using self-priming oligonucleotides containing the desired sequence. For the expression of clostridial neurotoxin fragments the codon bias and GC/AT base ratio was adjusted for ease of expression in *E. coli*. Fragments were cloned sequentially into pLitmus 38 (New England Biolabs, Inc., Beverly, MA) to assemble the entire gene. Constructs for expression were sub-cloned into pET28b (Novagen Inc., Madison, WI) replacing the *Eco*R1-*Hind*III fragment. The ligation reactions were transformed into *E. coli* DH5 $\alpha$  (Life Technologies Inc., Gaithersburg, MD). Plasmid DNA was amplified, purified and screened for the presence of the appropriate sequence (Ausubel *et al.* 1989, Current Protocols in Molecular Biology, John Wiley & Sons, New York). Gene constructions confirmed as possessing the correct sequences were then transformed into the expression host *E. coli* BL21 (DE3) (Studier & Moffatt 1986, *Journal of Molecular Biology*, 189: 113-130).

The recombinant constructs expressed in pET28 were produced with amino-terminal histidine (6 His) and T7 peptide tags allowing proteins to be purified by affinity chromatography on either a Ni<sup>2+</sup> charged column or an anti-T7 immunoaffinity column (Smith *et al.* 1988, *Journal of Biological Chemistry*, 263: 7211-7215). Incorporation of a factor X protease cleavage between the peptide tags and the expressed Mn-SOD Constructs allowed these to be removed after purification. Briefly, cultures of *E. coli* BL21 (DE3) pET28-Mn-SOD Construct were grown in Terrific broth-kanamycin (30  $\mu$ gml<sup>-1</sup>) to an OD<sub>600</sub> nm of 2.0, and protein expression was induced by the addition of 500  $\mu$ M IPTG for approximately 2 h. Cells were lysed by freeze/thaw followed by sonication, lysates cleared by centrifugation and supernatants loaded onto an anion exchange column (MonoQ™ column on

- 22 -

5 a Fast Protein Liquid Chromatography system; Pharmacia Biotech, Uppsala, Sweden). Eluted recombinant Mn-SOD Construct was then desalted and further purified by affinity chromatography on a chelating sepharose column charged with  $\text{Ni}^{2+}$  (Pharmacia Biotech, Uppsala, Sweden). After loading proteins onto the column and subsequent washing, the purified Construct was eluted with imidazole. All buffers used were as specified by the manufacturer.

10 A 'maltose binding protein' purification tag was also employed for the purification some batches of Mn-SOD Constructs. The use of this system is described in detail in New England Biolabs Instruction Manual "Protein Fusion and Purification System" (ver 3.02).

15 It would also be evident to anyone skilled in the art that other tags and protease cleavage site may also be incorporated into the sequence to facilitate purification as exemplified in Figure 3.

20 The amino sequences of several recombinant Mn-SOD constructs are shown in the sequence listing.

### Example 3.

#### Preparation of botulinum heavy chains by chemical methods.

25 The various serotypes of the clostridial neurotoxins may be prepared and purified from various toxigenic strains of *Clostridium botulinum* and *Clostridium tetani* by methods employing standard protein purification techniques as described previously (Shone and Tranter 1995, Current Topics in Microbiology, 194, 143-160; Springer). Samples of botulinum neurotoxin (1mg/ml) are dialysed against a buffer containing 50mM Tris-HCl pH 8.0, 1M NaCl and 2.5M urea for at least 4 hours at 4°C and then made 100mM with dithiothreitol and incubated for 16h at 22°C. The cloudy

30

- 23 -

5 solution which contains precipitated light chain is then centrifuged at 15000 x g for 2 minutes and the supernatant fluid containing the heavy chain retained and dialysed against 50mM Hepes pH 7.5 containing 0.2M NaCl and 5mM dithiothreitol for at least 4 hours at 4°C. The dialysed heavy chain is centrifuged at 15000 x g for 2 minutes and the supernatant retained and dialysed thoroughly against 50mM Hepes pH 7.5 buffer containing 0.2M NaCl and stored at -70°C. The latter procedure yields heavy chain >95% pure with a free cysteine residue which can be used for chemical coupling purposes. Biological (binding) activity of the heavy chain may be assayed as described in Example 5.

10 The heavy chains of the botulinum neurotoxins may also be produced by chromatography on QAE Sephadex as described by the methods in Shone and Tranter (1995) (Current Topics in Microbiology, 194, 143-160; Springer).

#### 15 Example 4

#### 20 Production of Mn-SOD constructs by chemical methods

25 *B. stearothermophilus* Mn-SOD fused to a mitochondrial leader sequence was purified as described in Example 1. The Mn-SOD was chemically modified by treatment with a 3-5 molar excess of N-succinimidyl 3-[2-pyridyldithio] propionate (SPDP) in 0.05M Hepes buffer pH 7.0 containing 0.1M NaCl for 60 min at 22°C. The excess SPDP was removed by dialysis against the same buffer at 4°C for 16h. The substituted SOD was then mixed in a 1:2.5 molar ratio with heavy chain purified from *Clostridium botulinum* type A neurotoxin purified as described in Example 3 and incubated at 4°C for 16h. During the incubation period the Mn-SOD was conjugated to the botulinum heavy chain fragment by free sulphydryl groups (see Figure 4). After incubation, the Mn-SOD-construct was purified by gel filtration chromatography on Sephadex G200.



- 24 -

Constructs of the invention may also be formed by the above method using polypeptides containing the translocation and binding domains that have been produced by recombinant technology as outlined in Example 2.

## Example 5.

### Assay of the biological activity of constructs - demonstration of high affinity binding to neuronal cells.

Clostridial neurotoxins may be labelled with 125-iodine using chloramine-T and its binding to various cells assessed by standard methods such as described in Evans *et al.* 1986, Eur J. Biochem., 154, 409 or Wadsworth *et al.* 1990, Biochem. J. 268, 123). In these experiments the ability of Mn-SOD constructs to compete with native clostridial neurotoxins for receptors present on neuronal cells or brain synaptosomes was assessed. All binding experiments were carried out in binding buffers. For the botulinum neurotoxins this buffer consisted of: 50mM HEPES pH 7.0, 30mM NaCl, 0.25% sucrose, 0.25% bovine serum albumin. For tetanus toxin, the binding buffer was: 0.05M tris-acetate pH 6.0 containing 0.6% bovine serum albumin. In a typical binding experiment the radiolabelled clostridial neurotoxin was held at a fixed concentration of between 1-20nM. Reaction mixtures were prepared by mixing the radiolabelled toxin with various concentrations of unlabelled neurotoxin or construct. The reaction mixture were then added to neuronal cells or rat brain synaptosomes and then incubated at 0-3°C for 2hr. After this period the neuronal cells of synaptosomes were washed twice with binding ice-cold binding buffer and the amount of labelled clostridial neurotoxin bound to cells or synaptosomes was assessed by  $\gamma$ -counting. In an experiment using an Mn-SOD construct which contained the binding domain from botulinum type A neurotoxin, the construct was found to compete with <sup>125</sup>I-labelled botulinum type A neurotoxin for neuronal cell receptors in a similar manner to unlabelled native botulinum type A neurotoxin. These data showed that

- 25 -

the construct had retained binding properties of the native neurotoxin.

### Example 6

#### 5      **Assay of the biological activity of constructs – measurement of the Mn-SOD activity.**

10      Mn-SOD activity in samples and constructs was measured by a modification (Brehm *et al.* (1991) Appl. Microbiol. Biotechnol., 36,358-363) of the procedure described by McCord and Fridovich (J. Biol. Chem. (1969), 244, 6049-6055). Aliquots (20 $\mu$ l) of samples or constructs containing Mn-SOD were added to 1ml of 0.05M potassium phosphate buffer pH 7.5 containing  $1 \times 10^{-4}$  M EDTA,  $2.5 \times 10^{-5}$  M ferricytochrome C and  $7 \times 10^{-3}$  M sodium xanthine in a thermostatted cuvette at 30°C. Sufficient xanthine oxidase was added to produce a rate of reduction of the ferricytochrome C at 550nm of approx. 0.1 absorbance units/minute in the absence of Mn-SOD. Under these conditions the amount Mn-SOD that was required to reduce the rate of reduction of ferricytochrome C by 50% was defined as one unit of activity.

20      Using such assays the Mn-SOD activity within constructs was assessed.

### Example 7

#### 25      **Demonstration of the targeting of Mn-SOD to the mitochondria of neuronal cells by constructs of the invention**

30      Mn-SOD construct containing the translocation and targeting domains derived from botulinum type A neurotoxin was incubated at various concentrations (0.01-10 $\mu$ M final concentration) with a neuroblastoma cell line NG108. Incubations were carried out over a 6h period or overnight at 37°C. In some experiments, construct radiolabelled with <sup>125</sup>I iodine was

- 26 -

used. After incubation with the construct cells, were removed from culture  
flasks by gentle scraping and centrifuged at 200 x g. Cells were then  
resuspended in breaking buffer (0.6M mannitol, 20mM hepes pH 7.4 and  
1mM phenylmethysulphonyl chloride) and homogenised in a Dounce  
homogeniser. The homogenate was centrifuged at 200 x g for 5min and  
then the supernatant fluid recovered and centrifuged at 8000 x g for  
10min. The 200 x g pellet (nuclear fraction) were pooled and resuspended  
in phosphate buffered saline. The 8000 x g pellets (mitochondrial fraction)  
were also pooled and resuspended in phosphate buffered saline. The  
supernatant fluid was saved and used to represent the cytosolic fraction.

Analysis of the sub-cellular distribution of Mn-SOD was carried out by  
Western blot analysis and, where radiolabelled construct was used, by  
analysis of the <sup>125</sup>I-labelled construct components by  $\gamma$ -counting and by  
autoradiography of cell fractions which had been separated by  
electrophoresis on SDS-polyacrylamide gels. For Western blot analysis,  
proteins in the cell fractions were separated by electrophoreses on SDS-  
polyacrylamide gels and then transferred to nitrocellulose membrane as  
described previously (Towbin *et al.* Proc.(1979) Natl. Acad. Sci. USA, 76,  
4350). The presence of Mn-SOD in protein bands on nitrocellulose  
membranes was assessed by incubation with rabbit anti- Mn-SOD antibody  
followed by washing and incubation with anti-rabbit peroxidase conjugate.  
Addition of peroxidase substrates (3,3',5,5'- tetramethyl benzidine and  
H<sub>2</sub>O<sub>2</sub>) allowed visualisation and quantitation of the Mn-SOD in the various  
sub-cellular protein fractions. An enhanced chemiluminescence system  
(Amersham International) was also used in some experiments to increase  
the sensitivity.

#### Example 8.

Formulation of the Mn-SOD construct for clinical use.

- 27 -

In a formulation of the Mn-SOD construct for clinical use, recombinant Mn-SOD construct would be prepared under current Good Manufacturing Procedures. The construct would be transferred, by dialysis, to a solution to give the product stability during freeze-drying. Such a formulation may contain Mn-SOD construct (10 mg/ml) in 5mM HEPES buffer (pH 7.2), 50mM NaCl, 1% lactose. The solution, after sterile filtration, would be aliquotted, freeze-dried and stored under nitrogen at -20°C.

#### Example 9.

##### Use of an Mn-SOD construct to treat stroke.

In a typical case of a middle aged or elderly man diagnosed as suffering from stroke, treatment with an Mn-SOD construct would begin immediately, ideally within 6 hours of the stroke occurring. Doses of the Mn-SOD construct (e.g. 100mg) reconstituted in a sterile saline solution would be administered intravenously. Further doses of the construct would be administered daily for 5-10 days. Such a patient would be expected to display reduced levels of ischaemia/reperfusion damage as assessed by magnetic resonance imaging compared to a similarly affected patient receiving no treatment. Relative improvements to muscle strength and co-ordination (MRC motor score) would be expected to be observed over the subsequent 12 month period.

#### Example 10.

##### Method to Demonstrate Neuroprotection of Cells by MnSOD-Heavy Chain Conjugates

Neuroblastoma cell line NG108-15 (*Nature* (1998) 336:p185) were seeded at a density of  $3 \times 10^4$  cells/ml in 96 well microtitre plates coated with poly-D-lysine. Plates were grown for 3 days at 37°C in a CO<sub>2</sub> incubator (5% CO<sub>2</sub>

- 28 -

95% air). MnSOD or leader-MnSOD conjugates were prepared with purified heavy chain of botulinum neurotoxin serotype A (BoNT/A HC) as described. The concentration of SOD was estimated and the conjugate diluted to give the specified amount of conjugate in a total volume of 200 $\mu$ l serum free medium. Conjugate was added to wells in the presence or absence of 56mM KCl, 2mM CaCl<sub>2</sub>. The cells were incubated with conjugate for 1 hour. The conjugate was replaced with either serum free medium or serum free medium containing 50 $\mu$ M duroquinone and incubated at 37°C for 4 hours in the CO<sub>2</sub> incubator to induce oxidative stress. The media was removed after 4 hours and replaced with the dye 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) at a final concentration of 0.25mg/ml in serum free medium and incubated for 2 hours (according to the method of Mattson, MP, *et al* (1995) *Methods in Cell Biology* 46:187-216) The conversion of MTT to formazan dye crystals has been shown to be related to mitochondrial respiratory chain activity (Musser, DA, and Oseroff, AR (1994) *Journal of Immunology* 59:621-626). MTT was removed and crystals solubilised with dimethylsulfoxide (DMSO). Absorbance at 570nm was measured using a Labsystems Multiskan Plus MkII spectrophotometer and the results shown in Figure 5.

Figure 5 shows that a construct of the invention was able to protect neuronal cells against the mitochondrial - focused oxidative stress produced by duroquinone.

#### Example 11.

##### Preparation and purification of recombinant MnSOD and leader sequence MnSOD

Standard molecular biology protocols were used for all genetic manipulations (Sambrook *et al* 1989, Molecular cloning; A laboratory manual. Second Edition, Cold Spring Harbor Laboratory Press, New York.).

- 29 -

5 The MnSOD gene from either *B. stearothermophilus* or *B. caldopenax* was amplified by PCR to engineer a *Bam*HI site (resulting in the replacement of nucleotides 1-15). A synthetic oligonucleotide corresponding to the mitochondrial leader sequence of human MnSOD (bases 1-81 of the human gene) was subcloned into the *Bam*HI site to generate leader-MnSOD. PCR was used to add a Factor Xa cleavage site immediately adjacent to the methionine at the start of the leader sequence. Similarly a Factor Xa cleavage site was engineered immediately adjacent to the methionine at the start of the native MnSOD gene. Constructs were sequenced to confirm the presence of the correct sequence. Constructs for expression were subcloned into the expression vector pET28a (Novagen Inc, Madison, WI) as an *Eco*RI fragment and the orientation of the fragments checked. Clones with confirmed sequences were used to transform expression host *E. coli* BL21 (DE3) (Studier and Moffatt 1986 *Journal of Molecular Biology* 189:113-130). Examples 2 and 3 above provide detailed methods.

20 The recombinant proteins expressed from pET28a contain amino-terminal histidine (6-His) and T7 peptide tags allowing proteins to be purified by affinity chromatography on either a  $\text{Cu}^{2+}$  charged metal chelate column or an anti-T7 immunoaffinity column. Incorporation of the Factor Xa site between the peptide tags and the start of either MnSOD or leader MnSOD allows the precise removal of the peptide tags after purification. Cultures of *E. coli* BL21(DE3) pET28a-MnSOD or BL21(DE3) pET28a-leader-MnSOD were grown in Terrific Broth containing 30  $\mu\text{g/ml}$  kanamycin and 0.5% (w/v) glucose to an  $\text{OD}_{600}$  of 2.0 and protein expression was induced with 500  $\mu\text{M}$  IPTG for 2 hours. Cells were lysed by sonication, cell debris pelleted by centrifugation and the supernatant loaded onto a metal chelate column charged with  $\text{Cu}^{2+}$  (Amersham-Pharmacia Biotech, Uppsala, Sweden). After loading proteins on the column and washing, proteins were eluted using imidazole. All buffers were used as specified by manufacturers. Factor Xa cleavage of the eluted protein was carried out according to manufacturers instructions.

- 30 -

The invention thus provides constructs and methods for delivery of SOD to neuronal cells.

FOUO - 030799

FIG. 1






FIG. 2





2/4

FIG. 3



3/4





FIG. 4

**FIG. 5** Demonstration of protective effects of HC/A Mnsod on NG108 cells subjected to oxidative stress by the addition of 50 $\mu$ M duroquinone for 4 hours.



Variable	Mean	SD	Min	Max
Age	38.5	10.5	25	55
Gender	Male			
Marital status	Married			
Education	High school			
Occupation	Manager			
Income	10,000	5,000	5,000	20,000
Health status	Good			
Exercise frequency	3 times/week			
Stress level	Low			
Sleep quality	Good			
Dietary habits	Healthy			
Alcohol consumption	Low			
Tobacco use	None			
Family size	3			
Work hours	40			
Commuting time	30			
Living space	100			
Neighborhood safety	High			
Access to green spaces	Yes			
Public transportation	Good			
Local economy	Stable			
Community support	Strong			
Healthcare access	Good			
Environmental quality	High			
Crime rate	Low			
Local government	Effective			
Local culture	Rich			
Local history	Long			
Local industry	Diverse			
Local infrastructure	Modern			
Local services	Complete			
Local environment	Healthy			
Local climate	Temperate			
Local vegetation	Lush			
Local wildlife	Diverse			
Local water quality	Good			
Local air quality	Good			
Local noise level	Low			
Local traffic volume	Low			
Local parking availability	Good			
Local shopping options	Many			
Local dining options	Many			
Local entertainment options	Many			
Local cultural events	Many			
Local historical sites	Many			
Local parks and recreation	Many			
Local sports facilities	Many			
Local medical facilities	Many			
Local educational facilities	Many			
Local government services	Many			
Local public services	Many			
Local private services	Many			
Local social services	Many			
Local religious services	Many			
Local cultural services	Many			
Local historical services	Many			
Local environmental services	Many			
Local water services	Many			
Local air services	Many			
Local noise services	Many			
Local traffic services	Many			
Local parking services	Many			
Local shopping services	Many			
Local dining services	Many			
Local entertainment services	Many			
Local cultural services	Many			
Local historical services	Many			
Local environmental services	Many			
Local water services	Many			
Local air services	Many			
Local noise services	Many			
Local traffic services	Many			
Local parking services	Many			
Local shopping services	Many			
Local dining services	Many			
Local entertainment services	Many			
Local cultural services	Many			
Local historical services	Many			
Local environmental services	Many			
Local water services	Many			
Local air services	Many			
Local noise services	Many			
Local traffic services	Many			
Local parking services	Many			
Local shopping services	Many			
Local dining services	Many			
Local entertainment services	Many			
Local cultural services	Many			
Local historical services	Many			
Local environmental services	Many			
Local water services	Many			
Local air services	Many			
Local noise services	Many			
Local traffic services	Many			
Local parking services	Many			
Local shopping services	Many			
Local dining services	Many			
Local entertainment services	Many			
Local cultural services	Many			
Local historical services	Many			
Local environmental services	Many			
Local water services	Many			
Local air services	Many			
Local noise services	Many			
Local traffic services	Many			
Local parking services	Many		</	

As a below named inventor, I hereby declare that:

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter that is claimed and for which a patent is sought on the invention entitled "Delivery of Superoxide Dismutase to Neuronal Cells", the specification of which is attached hereto unless the following box is checked:

☐ was filed on \_\_\_\_\_;  
as United States Application Number or PCT International Application Number \_\_\_\_\_, and  
was amended on \_\_\_\_\_ (if applicable)

I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application, which designated at least one country other than the United States listed below, and have also identified below any foreign application for patent or inventor's certificate, or PCT international application having a filing date before that of the application on which priority is claimed

Prior Foreign Application(s)			Priority	Claimed
<u>9824282.9</u>	<u>United Kingdom</u>	<u>5 November 1998</u>	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
(Application No.)	(Country)	(Day/Month/Year Filed)		
<u>                    </u>	<u>                    </u>	<u>                    </u>	<input type="checkbox"/> Yes	<input type="checkbox"/> No
(Application No.)	(Country)	(Day/Month/Year Filed)		

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

(Application No.) (Filing Date)

(Application No.) \_\_\_\_\_ (Filing Date) \_\_\_\_\_

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or under § 365(c) of any PCT international application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56 that became available between the filing date of the prior application and the national or PCT international filing date of this application.

PCT/GB99/03699 (Application No.)	5 November 1999 (Filing Date)	Pending (Status - patented, pending, abandoned)
(Application No.)	(Filing Date)	(Status - patented, pending, abandoned)

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

12 Robert G. Sterne, Esq., Reg. No. 28,912; Edward J. Kessler, Esq., Reg. No. 25,688; Jorge A. Goldstein, Esq., Reg. No. 29,021; Samuel L. Fox, Esq., Reg. No. 30,353; David K S Cornwell, Esq., Reg. No. 31,944; Robert W. Esmond, Esq., Reg. No. 32,893; Tracy-Gene G. Durkin, Esq., Reg. No. 32,831; Michele A. Cimbala, Esq., Reg. No. 33,851; Michael B. Ray, Esq., Reg. No. 33,997; Robert E. Sokohl, Esq., Reg. No. 36,013; Eric K. Steffe, Esq., Reg. No. 36,688; and Michael Q. Lee, Esq., Reg. No. 35,239.

I hereby authorize the U.S. attorneys named herein to accept and follow instructions from Mathys & Squire, 100 Gray's Inn Road, London WC1X 8AL, ENGLAND, as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. attorneys and me. In the event of a change in the persons from whom instructions may be taken, the U.S. attorneys named herein will be so notified.

Send Correspondence to:


STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.  
1100 New York Avenue, N.W., Suite 600  
Washington, D.C. 20005-3934


Direct Telephone Calls to:

(202) 371-2600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor	Clifford Charles Shone
Signature of sole or first inventor	Date 30/07/01
Residence	United Kingdom - Salisbury, Wiltshire
Citizenship	British
Post Office Address	c/o Microbiological Research Authority, CAMR, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom
Full name of second inventor	John Mark Sutton
Signature of second inventor	Date 30/07/01
Residence	United Kingdom - Salisbury, Wiltshire
Citizenship	British
Post Office Address	c/o Microbiological Research Authority, CAMR, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom

Full name of third inventor	<u>Bassam Hallis</u>	Date	30/07/01
Signature of third inventor			
Residence	United Kingdom - <u>Salisbury</u> , Wiltshire		
Citizenship	United Kingdom		
Post Office Address	c/o Microbiological Research Authority, CAMR, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom		

Full name of fourth inventor	<u>Nigel Silman</u>		
Signature of fourth inventor		Date	30/07/01
Residence	United Kingdom – <u>Salisbury, Wiltshire</u>		
Citizenship	United Kingdom		
Post Office Address	c/o Microbiological Research Authority, CAMR, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom		
Full name of fifth inventor			
Signature of fifth inventor			Date
Residence			
Citizenship			
Post Office Address			

H:\WPDOCS\GWS\FORMS\USFORMS\20994DEC  
SKGE Rev. 9/2/98 mac

(Supply similar information and signature for subsequent joint inventors, if any)

## POWER OF ATTORNEY FROM ASSIGNEE WITH DELEGATION

Microbiological Research Authority, having a principal place of business at CAMR, Porton Down, Salisbury, Wiltshire SP4 0JG, Great Britain, is assignee of the entire right, title, and interest for the United States of America (as defined in 35 U.S.C. §100), by reason of an Assignment to the Assignee executed on 30 July 2001 of an invention known as Delivery of Superoxide Dismutase to Neuronal Cells (Attorney Docket No. 1581.0800000), which is disclosed and claimed in a patent application of the same title by the inventor(s) Clifford Charles Shone, John Mark Sutton, Bassam Hallis and Nigel Silman (said application filed on 09/031,050 at the U.S. Patent and Trademark Office, having Application Number 1-A.Filing Date: Nov. 5, 1999.

Reg No.  
32,893

The Assignee hereby appoints the following U.S. attorneys to prosecute this application and any continuation, divisional, continuation-in-part, or reissue application thereof, and to transact all business in the U.S. Patent and Trademark Office connected therewith: Robert Greene Sterne, Registration No 28,912; Edward J Kessler, Registration No 25,688; Jorge A. Goldstein, Registration No. 29,021; Samuel L Fox, Registration No 30,353; David K.S. Cornwell, Registration No 31,944; Robert W Esmond, Registration No. 32,893; Tracy-Gene G. Durkin, Registration No. 32,831; Michele A. Cimbala, Registration No. 33,851, Michael B. Ray, Registration No. 33,997; Robert E. Sokohl, Registration No. 36,013; Eric K. Steffe, Registration No. 36,688; Michael Q Lee, Registration No. 35,239; and Steven R. Ludwig, Registration No. 36,203. The Assignee hereby grants said attorneys the power to insert on this Power of Attorney any further identification that may be necessary or desirable in order to comply with the rules of the U.S. Patent and Trademark Office.

The Assignee hereby authorizes the U.S. attorneys named herein to accept and follow instructions from Mathys & Squire, 100 Gray's Inn Road, London WC1X 8AL, Great Britain as to any action to be taken in the U S Patent and Trademark Office regarding this application without direct communication between the U.S. attorneys and the Assignee. In the event of a change in the persons from whom instructions may be taken, the U S attorneys named herein will be so notified by the Assignee.

Send correspondence to:

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.  
1100 New York Avenue, N.W.  
Suite 600  
Washington, D.C. 20005-3934  
U.S.A.

Direct phone calls to 202-371-2600.

FOR: Microbiological Research Authority  
SIGNATURE: [Signature]  
BY: [Signature]  
TITLE: SECRETARY MRA (CAME)  
DATE: 31 July 2001